Formulary Adherence Checklist for NICE Technology Appraisals About Medicines

Milton Keynes Health Economy



Developed for the NHS by: East and South East England Specialist Pharmacy Services

This spreadsheet is updated monthly and enables self-audit of a medicines formulary for adherence to current NICE Technology Appraisals. Version 20 January 2018 All guidelines refer to adults unless indicated. No copyright is asserted on this material if used for non-commercial purposes within the NHS.

Technology appraisal (TA) Titles are hyperlinks to full guidance	Date of TA Release	Availability of medicine for NHS patients with this medical condition, as indicated by NICE							
			Yes (mark 'x' if applicable)	N/A (mark 'x' if applicable)	Date of local decision (DD/MM/YY)	Time to implement (days)	Notes (e.g. rationale, method of making available)		
2017/2018 Golimumab for treating non- radiographic axial spondyloarthritis (TA497)		Golimumab (Simponi®) is recommended, within its marketing authorisation, as an option for treating severe non-radiographic axial spondyloarthritis in adults whose disease has responded inadequately to, or who cannot tolerate, nonsteroidal anti-inflammatory drugs.	х		24/01/2018	14	CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements		
Ribociclib with an aromatase inhibitor for previously untreated, hormone receptorpositive, HER2-negative, locally advanced or metastatic breast cancer (TA496)	20/12/2017	Ribociclib (Kisqali), with an aromatase inhibitor, is recommended within its marketing authorisation, as an option for treating hormone receptor-positive, human epidermal growth factor receptor 2-negative, locally advanced or metastatic breast cancer as initial endocrine-based therapy in adults	x		24/01/2018	35	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements		
Palbociclib with an aromatase inhibitor for previously untreated, hormone receptorpositive, HER2-negative, locally advanced or metastatic breast cancer (TA495)		Palbociclib (Ibrance), with an aromatase inhibitor, is recommended within its marketing authorisation, as an option for treating hormone receptor-positive, human epidermal growth factor receptor 2-negative, locally advanced or metastatic breast cancer as initial endocrine-based therapy in adults	x		24/01/2018	35	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements		
Naltrexone–bupropion for managing overweight and obesity (TA494)	12/12/2017	Naltrexone-bupropion (Mysimba) is not recommended within its marketing authorisation for managing overweight and obesity in adults alongside a reduced-calorie diet and increased physical activity	х		24/01/2018	43	Not recommended		
Cladribine tablets for treating relapsing–remitting multiple sclerosis (TA493)	06/12/2017	Cladribine tablets (Mavenclad) are recommended as an option for treating highly active multiple sclerosis in adults	х		24/01/2018		NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements		

Atezolizumab for untreated locally advanced or metastatic urothelial cancer when cisplatin is unsuitable (TA492)	06/12/2017	Atezolizumab (Tecentriq) is recommended for use within the Cancer Drugs Fund as an option for untreated locally advanced or metastatic urothelial carcinoma in adults, for whom cisplatin-based chemotherapy is unsuitable Ibrutinib (Imbruvica) is recommended for use in	х	24/01/2018	49 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements 63 NHSE commissioned. Approved for addition to the formulary and to be used in line with
<u>Ibrutinib for treating</u> <u>Waldenstrom's</u> <u>macroglobulinaemia (TA491)</u>	22/11/2017	the Cancer Drugs Fund as an option for treating Waldenstrom's macroglobulinaemia in adults who have had at least 1 prior therapy	x	24/01/2018	NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements
Nivolumab for treating squamous cell carcinoma of the head and neck after platinum-based chemotherapy (TA490)	22/11/2017	Nivolumab (Opdivo) is recommended for use within the Cancer Drugs Fund as an option for treating squamous cell carcinoma of the head and neck in adults whose disease has progressed on platinum-based chemotherapy	х	24/01/2018	63 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements
Vismodegib for treating basal cell carcinoma (TA489)	22/11/2017	Vismodegib (Erivedge) is not recommended within its marketing authorisation for treating symptomatic metastatic basal cell carcinoma, or locally advanced basal cell carcinoma that is inappropriate for surgery or radiotherapy	x	24/01/2018	63 Not recommended
Regorafenib for previously treated unresectable or metastatic gastrointestinal stromal tumours (TA488)	15/11/2017	Regorafenib (Stivarga) is recommended as an option for treating unresectable or metastatic gastrointestinal stromal tumours in adults whose disease has progressed on, or who are intolerant to, prior treatment with imatinib and sunitinib	x	24/01/2018	70 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements
Venetoclax for treating chronic lymphocytic leukaemia (TA487)	01/10/2017	Venetoclax (Venclyxto®) is recommended for use within the Cancer Drugs Fund, within its marketing authorisation, as an option for treating chronic lymphocytic leukaemia	x	29/11/2017	59 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements
Aflibercept for treating choroidal neovascularisation (TA486)	01/10/2017	Aflibercept (Eylea®) is recommended, within its marketing authorisation, as an option for treating visual impairment because of myopic choroidal neovascularisation in adults	х	29/11/2017	59 CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Sarilumab for moderate to severe rheumatoid arthritis (TA485)	01/10/2017	Sarilumab (Kevzara®) with methotrexate, is recommended as an option for treating active rheumatoid arthritis in adults whose disease has responded inadequately to intensive therapy with a combination of conventional diseasemodifying anti-rheumatic drugs (DMARDs)	х	29/11/2017	59 CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Nivolumab for previously treated non-squamous non- small-cell lung cancer (TA484)	01/10/2017	Nivolumab (Opdivo®) is recommended for use within the Cancer Drugs Fund as an option for treating locally advanced or metastatic non-squamous non-small-cell lung cancer in adults after chemotherapy	х	29/11/2017	59 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements

Nivolumab for previously treated squamous non-small- cell lung cancer (TA483)	01/10/2017	within the Cancer Drugs Fund as an option for treating locally advanced or metastatic squamous non-small-cell lung cancer in adults after chemotherapy	x	29/11/2017		NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements
Immunosuppressive therapy for kidney transplant in children and young people (TA482)	01/10/2017	Immunosuppressive therapy for kidney transplant in children and young people	x	29/11/2017	59	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Immunosuppressive therapy for kidney transplant in adults (TA481)	01/10/2017	Immunosuppressive therapy for kidney transplant in adults	х	29/11/2017	59	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Tofacitinib for moderate to severe rheumatoid arthritis (TA480)	01/10/2017	Tofacitinib (Xeljanz®) is recommended as an option for treating active rheumatoid arthritis in adults whose disease has responded inadequately to intensive therapy with a combination of conventional disease-modifying anti-rheumatic drugs (DMARDs)	x	29/11/2017		CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Reslizumab for treating severe eosinophilic asthma (TA479)	01/10/2017	Reslizumab (Cinqaero®) is recommended as an option for the treatment of severe eosinophilic asthma that is inadequately controlled in adults despite maintenance therapy with high-dose inhaled corticosteroids plus another drug	x	29/11/2017	59	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Brentuximab vedotin for treating relapsed or refractory systemic anaplastic large cell lymphoma (TA478)	01/10/2017	Brentuximab vedotin (Adcetris®) is recommended as an option for treating relapsed or refractory systemic anaplastic large cell lymphoma in adults	x	29/11/2017	59	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements and the National Cancer Drugs Fund requirements
Autologous chondrocyte implantation for treating symptomatic articular cartilage defects of the knee (TA477)	01/10/2017	Autologous chondrocyte implantation (ACI) ChondroCelect® is recommended as an option for treating symptomatic articular cartilage defects of the knee	х	29/11/2017	59	NICE Appraisal N/A at MK
Paclitaxel as albumin-bound nanoparticles with gemcitabine for untreated metastatic pancreatic cancer (TA476)	06/09/2017	Paclitaxel as albumin-bound nanoparticles (nab-paclitaxel) with gemcitabine is recommended as an option for untreated metastatic adenocarcinoma of the pancreas in	х	27/09/2017	21	Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Dimethyl fumarate for treating moderate to severe plaque psoriasis (TA475)	06/09/2017	Dimethyl fumarate is recommended as an option for treating plaque psoriasis in adults	х	29/11/2017	84	Confirmed place in therapy with dermatologists. CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements.
Sorafenib for treating advanced hepatocellular carcinoma (TA474)		Sorafenib is recommended as an option for treating advanced hepatocellular carcinoma only for people with Child-Pugh grade A liver impairment	х	27/09/2017		Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Cetuximab for treating recurrent or metastatic squamous cell cancer of the head and neck (TA473)	31/08/2017	Cetuximab in combination with platinum-based chemotherapy is recommended as an option for treating recurrent or metastatic squamous cell cancer of the head and neck in adults	x	27/09/2017	27	CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements.

Obinutuzumab with	30/08/2017	Obinutuzumab in combination with		27/09/2017	29 Ann	proved for addition to the formulary and to be used in line with NICE guidance and
bendamustine for treating	30/08/2017	bendamustine followed by obinutuzumab		27/09/2017		nmissioning statements
		-			Con	minissioning statements
follicular lymphoma refractory to		maintenance is recommended for use within the				
<u>rituximab (TA472)</u>		Cancer Drugs Fund as an option for treating	x			
		adults with follicular lymphoma that did not				
		respond or progressed during or up to 6 months				
		after treatment with rituximab or a rituximab-				
		containing regimen				
Eluxadoline for treating irritable	30/08/2017	Eluxadoline is recommended as an option for		27/09/2017	28 App	proved for addition to the formulary and to be used in line with NICE guidance with
bowel syndrome with diarrhoea		treating irritable bowel syndrome with diarrhoea	x		Sec	ondary Care Initiation and stabilisation including one month review.
(TA471)		in adults	^			
Ofatumumab with chemotherapy	23/08/2017	Ofatumumab with chemotherapy - was unable		27/09/2017	35 Ter	minated Appraisal by NICE
for treating chronic lymphocytic		to make a recommendation about the use in the				
leukaemia (terminated appraisal)		NHS of ofatumumab with chemotherapy for				
<u>(TA470)</u>		treating chronic lymphocytic leukaemia because	х			
		no evidence submission was received from				
		Novartis Pharmaceuticals UK				
Idelalisib with ofatumumab for	23/08/2017	NICE is unable to make a recommendation about		27/09/2017	2E Tori	minated Appraisal by NICE
treating chronic lymphocytic	23/00/201/	the use in the NHS of idelalisib with ofatumumab		21/03/2011	35 161	minated Appraisal by NICL
			X			
leukaemia (terminated appraisal)		for treating chronic lymphocytic leukaemia				
(TA469)	22/09/2017	because no evidence submission was received		27/00/2017	25 Tor:	mineted Appraisal by NICE
Methylnaltrexone bromide for	23/08/2017	Methylnaltrexone bromide - was unable to make		27/09/2017	35 Teri	minated Appraisal by NICE
treating opioid-induced		a recommendation about the use in the NHS of				
constipation (terminated		methylnaltrexone bromide for treating opioid-	х			
appraisal) (TA468)		induced constipation because no evidence				
		submission was received from Swedish Orphan				
		Biovitrum Ltd				
Holoclar for treating limbal stem	16/08/2017	Holoclar (ex vivo expanded autologous human		27/09/2017	42 App	proved for addition to the formulary and to be used in line with NICE guidance and
cell deficiency after eye burns		corneal epithelial cells containing stem cells) is			com	nmissioning statements
(TA467)		recommended as an option in people with	x			
<u> </u>		moderate to severe limbal stem cell deficiency				
		after eve burns				
Baricitinib for moderate to severe	09/08/2017			27/09/2017	49 App	proved for addition to the formulary and to be used in line with NICE guidance and
rheumatoid arthritis (TA466)		Baricitinib, with methotrexate is recommended			com	nmissioning statements
		as an option for treating active rheumatoid				
		arthritis in adults whose disease has responded	x			
		inadequately to intensive therapy with a				
		combination of conventional disease-modifying				
		antirheumatic drugs (DMARDs),				
Olaratumab in combination with	09/08/2017	Olaratumab, in combination with doxorubicin,		27/09/2017	49 App	proved for addition to the formulary and to be used in line with NICE guidance and
doxorubicin for treating advanced		is recommended for use within the Cancer Drugs	,,			nmissioning statements
soft tissue sarcoma (TA465)		Fund as an option for advanced soft tissue	x			-
		·				
Bisphosphonates for treating	09/08/2017	sarcoma in adults Oral bisphosphonates (alendronic acid,		27/09/2017	49 Apr	proved for addition to the formulary and to be used in line with NICE guidance and
osteoporosis (TA464)		ibandronic acid and risedronate sodium) are		, ,		nmissioning statements
		recommended as options for treating	х		3011	5
		osteoporosis in adults				
-				 		

Cabozantinib for previously treated advanced renal cell carcinoma (TA463)	09/08/2017	Cabozantinib is recommended, within its marketing authorisation, as an option for treating advanced renal cell carcinoma in adults after vascular endothelial growth factor (VEGF)-targeted therapy	х	27/09/2017	49 Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Nivolumab for treating relapsed or refractory classical Hodgkin lymphoma (TA462)	26/07/2017	Nivolumab is recommended, within its marketing authorisation, as an option for treating relapsed or refractory classical Hodgkin lymphoma in adults after autologous stem cell transplant and treatment with brentuximab vedotin, when the company provides nivolumab with the discount agreed in the patient access scheme.	х	27/09/2017	63 Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Roflumilast for treating chronic obstructive pulmonary disease (TA461)	26/07/2017	Roflumilast, as an add-on to bronchodilator therapy, is recommended as an option for treating severe chronic obstructive pulmonary disease in adults with chronic bronchitis	х	27/09/2017	63 Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Adalimumab and dexamethasone for treating non-infectious uveitis (TA460)	26/07/2017	Adalimumab is recommended as an option for treating non-infectious uveitis in the posterior segment of the eye in adults with inadequate response to corticosteroids. Dexamethasone (Ozurdex®) intravitreal implant is recommended as an option for treating non-infectious uveitis in the posterior segment of the eye in adults	x	27/09/2017	63 Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Collagenase clostridium histolyticum for treating Dupuytren's contracture (TA459)	26/07/2017	Collagenase clostridium histolyticum for people who meet the inclusion criteria for the ongoing clinical trial (HTA-15/102/04), comparing collagenase clostridium histolyticum (CCH) with limited fasciectomy, are encouraged to participate in the study. For people not taking part in the ongoing clinical trial, CCH is recommended as an option for treating Dupuytren's contracture with a palpable cord in adults	x	27/09/2017	63 Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Trastuzumab emtansine for treating HER2-positive advanced breast cancer after trastuzumab and a taxane (TA458)	19/07/2017		x	27/09/2017	Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Carfilzomib for previously treated multiple myeloma (TA457)	19/07/2017	Carfilzomib in combination with dexamethasone is recommended as an option for treating multiple myeloma in adults	х	27/09/2017	70 Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements

Ustekinumab for moderately to severely active Crohn's disease after previous treatment (TA456)	12/07/2017	Ustekinumab is recommended, within its marketing authorisation, as an option for treating moderately to severely active Crohn's disease, that is, for adults who have had an inadequate response with, lost response to, or were intolerant to either conventional therapy or a TNF-alpha inhibitor or have medical contraindications to such therapies.	х	27/09/2017		Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Adalimumab, etanercept and ustekinumab for treating plaque psoriasis in children and young people (TA455)	12/07/2017	Adalimumab, Etanercept and Ustekinumab is recommended as an option for treating plaque psoriasis in children and young people aged 4 years or older	х	27/09/2017		Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Daratumumab with lenalidomide and dexamethasone for treating relapsed or refractory multiple myeloma (terminated appraisal) (TA454)	05/07/2017	Daratumumab - was unable to make a recommendation about the use in the NHS of daratumumab, with lenalidomide and dexamethasone, for treating relapsed or refractory multiple myeloma because no evidence submission was received from Janssen-Cilag	х	26/07/2017	21	Terminated Appraisal by NICE
Bortezomib for treating multiple myeloma after second or subsequent relapse (terminated appraisal) (TA453)	05/07/2017	Bortezomib - was unable to make a recommendation about the use in the NHS of bortezomib for treating multiple myeloma after second or subsequent relapse because no evidence submission was received from Janssen-Cilag	x	26/07/2017	21	Terminated Appraisal by NICE
Ibrutinib for untreated chronic lymphocytic leukaemia without a 17p deletion or TP53 mutation (terminated appraisal) (TA452)	05/07/2017	Ibrutinib - was unable to make a recommendation about the use in the NHS of ibrutinib for untreated chronic lymphocytic leukaemia without a 17p deletion or TP53 mutation because no evidence submission was received from Janssen–Cilag	x	26/07/2017	21	Terminated Appraisal by NICE
Ponatinib for treating chronic myeloid leukaemia and acute lymphoblastic leukaemia (TA451)	28/06/2017	Ponatinib (Iclusig) is recommended, within its marketing authorisation, as an option for treating chronic-, accelerated- or blast-phase chronic myeloid leukaemia in adults	x	26/07/2017		NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Blinatumomab for previously treated Philadelphia-chromosome negative acute lymphoblastic leukaemia (TA450)	28/06/2017	Blinatumomab (Blincyto) is recommended within its marketing authorisation as an option for treating Philadelphia-chromosome-negative relapsed or refractory precursor B-cell acute lymphoblastic leukaemia in adults, only if the company provides it with the discount agreed in the patient access scheme.	x	26/07/2017		NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Everolimus and sunitinib for treating unresectable or metastatic neuroendocrine tumours in people with progressive disease (TA449)	28/06/2017	Everolimus (Afinitor) and Sunitinib (Sutent) are recommended, within their marketing authorisations, as options for treating well- or moderately differentiated unresectable or metastatic neuroendocrine tumours (NETs) of pancreatic origin in adults with progressive disease.	х	26/07/2017		NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements

Etelcalcetide for treating secondary hyperparathyroidism (TA448) Pembrolizumab for untreated PD- L1-positive metastatic non-small- cell lung cancer (TA447)	28/06/2017 28/06/2017	Etelcalcetide (Parsabiv) is recommended as an option for treating secondary hyperparathyroidism in adults with chronic kidney disease on haemodialysis Pembrolizumab (Keytruda) is recommended for use within the Cancer Drugs Fund as an option for untreated PD-L1-positive metastatic non-	x		26/07/2017	28 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements 28 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Brentuximab vedotin for treating CD30-positive Hodgkin lymphoma (TA446)	28/06/2017	small-cell lung cancer in adults Brentuximab vedotin (Adcetris) is recommended as an option for treating CD30-positive Hodgkin lymphoma in adults	x		26/07/2017	28 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Certolizumab pegol and secukinumab for treating active psoriatic arthritis after inadequate response to DMARDs (TA445)	24/05/2017	Certolizumab pegol (Cimzia) alone, or in combination with methotrexate is recommended as an option for treating active psoriatic arthritis in adults	x		26/07/2017	63 CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements.
Afatinib for treating advanced squamous non-small-cell lung cancer after platinum-based chemotherapy (terminated appraisal) (TA444)	22/03/2017	Tenofovir alafenamide - was unable to make a recommendation about the use in the NHS of afatinib for treating locally advanced or metastatic squamous non-small-cell lung cancer after platinum-based chemotherapy because no evidence submission was received from Boehringer Ingelheim.	x		24/05/2017	63 Terminated Appraisal by NICE
Obeticholic acid for treating primary biliary cholangitis (TA443)	26/04/2017	Obeticholic acid (Ocaliva) is recommended, within its marketing authorisation, as an option for treating primary biliary cholangitis in combination with ursodeoxycholic acid for people whose disease has responded inadequately to ursodeoxycholic acid or as monotherapy for people who cannot tolerate ursodeoxycholic acid. Obeticholic acid is recommended only if the company provides it with the discount agreed in the patient access	x		24/05/2017	NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Ixekizumab for treating moderate to severe plaque psoriasis (TA442)	26/04/2017	scheme. Ixekizumab (Taltz) is recommended as an option for treating plaque psoriasis in adults	х		24/05/2017	28 CCG commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements.
Daclizumab for treating relapsing—remitting multiple sclerosis (TA441)	26/04/2017	Daclizumab (Zinbryta) is recommended as an option for treating multiple sclerosis in adults	х		24/05/2017	28 NHSE commissioned. Approved for addition to the formulary and to be used in line with NICE guidance and commissioning statements
Pegylated liposomal irinotecan for treating pancreatic cancer after gemcitabine (TA440)	26/04/2017	Pegylated liposomal irinotecan (Onivyde), in combination with 5-fluorouracil and leucovorin, is not recommended, within its marketing authorisation, for treating metastatic adenocarcinoma of the pancreas in adults whose disease has progressed after gemcitabine-based therapy.	x		24/05/2017	28 Not recommended
			58	0		

		% "Yes"	% "N/A"	-	Average implement time (days)	
Adherence statistics for 2017-18		100%	0%		47	i i